
# **Emergency Care**

THIRTEENTH EDITION



CHAPTER 1

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

Respiratory Emergencies

# Multimedia Directory

| Slide 80 | Chronic Obstructive Pulmonary Diseases Video |
|----------|--|
| Slide 81 | Spontaneous Pneumothorax Animation           |
| Slide 90 | Metered-Dose Inhaler Video                   |

#### Topics

- Respiration
- Breathing Difficulty
- Respiratory Conditions
- The Prescribed Inhaler
- The Small-Volume Nebulizer


Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

#### Respiration

# Respiratory Anatomy and Physiology

- Diaphragm is muscular structure that separates the chest cavity from the abdominal cavity.
- During normal respiratory cycle, diaphragm and other parts of body work together to inhale and exhale.

# Respiratory Anatomy and Physiology



RELAXED

The process of respiration.

# Respiratory Anatomy and Physiology

- Inspiration
  - Active process
    - Uses muscle contraction to increase size of chest cavity
  - Intercostal muscles and diaphragm contract.
  - Diaphragm lowers; ribs move upward and outward.
  - Air is pulled into lungs.

continued on next slide

# Respiratory Anatomy and Physiology

- Expiration
  - Passive process
  - Rib muscles and diaphragm relax
  - Size of chest cavity decreases
  - Air flows out of lungs

#### Adequate Breathing

- Breathing sufficient to support life
- Signs

ALWAYS LEARNING

- No obvious distress
- Ability to speak in full sentences without having to catch his breath
- Normal color, mental status, and orientation

continued on next slide

#### Adequate Breathing

- May be determined by observing rate, rhythm, quality
  - 12 to 20 breaths/minute for adult
  - 15 to 30 breaths/minute for child
  - 25 to 50 breaths/minute for infant
  - Rhythm usually regular
  - Breath sounds normally present and equal

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

#### Inadequate Breathing

- Breathing not sufficient to support life.
- Signs
  - Rate out of normal range
  - Irregular rhythm
  - Diminished or absent lung sounds
  - Poor tidal volume

#### Pediatric Note

- Structure of an infant's and child's airway differs from that of an adult.
  - Smaller airway easily obstructed
  - Proportionately larger tongues
  - Smaller, softer, more flexible trachea
  - Less developed, less rigid cricoid cartilage
  - Heavy dependence on diaphragm for respiration

continued on next slide

#### Pediatric Note

- Signs of inadequate breathing in infants and children
  - Nasal flaring
  - Grunting
  - Seesaw breathing
  - Retractions

#### Patient Care

- Inadequate Breathing
  - Assisted ventilation with supplemental oxygen
    - Pocket face mask with supplemental oxygen
    - Two-rescuer bag-valve mask with supplemental oxygen

continued on next slide

ALWAYS LEARNING

#### Patient Care

- Inadequate Breathing
  - Assisted ventilation with supplemental oxygen
    - Flow-restricted, oxygen-powered ventilation device (FROPVD)
    - One-rescuer bag-valve mask with supplemental oxygen

#### Adequate and Inadequate **Artificial Ventilation**

- Chest rise and fall should be visible with each breath.
- Adequate artificial ventilation rates
  - 12 breaths per minute for adults
  - 20 breaths per minute for infants and children

continued on next slide

ALWAYS LEARNING


#### Adequate and Inadequate Artificial Ventilation

- Increasing pulse rates can indicate inadequate artificial ventilation in adults.
- Decreasing pulse rates can indicate inadequate artificial ventilation in pediatric patients.

#### Think About It

- How might you recognize the progression from adequate breathing to inadequate breathing in the assessment of your patient?
- How might your patient change during this transition?

- Patient's subjective perception
- Feeling of labored, or difficult, breathing
- Amount of distress felt may or may not reflect actual severity of condition.



1. Assess the patient and ensure that he meets the criteria for CPAP.

- Onset
  - When did it begin?
- Provocation
  - What were you doing when this came on?
- Quality

ALWAYS LEARNING

Do you have a cough? Are you bringing anything up with it?

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

continued on next slide

Radiation

ALWAYS LEARNING

Do you have pain or discomfort anywhere else in your body? Does it seem to spread to any other part of your body?

continued on next slide

All Rights Reserved

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

- Severity
  - On a scale of 1 to 10, how bad is your breathing trouble?
- Time

ALWAYS LEARNING

How long have you had this feeling?

continued on next slide

Observing

ALWAYS LEARNING

- Altered mental status
- Unusual anatomy
  - Barrel chest
- Patient's position
  - Tripod position
  - Sitting with feet dangling, leaning forward

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.


continued on next slide

- Observing
  - Work of breathing
    - Retractions
    - Use of accessory muscles
    - Flared nostrils
    - Pursed lips
    - Number of words patient can say without stopping

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

continued on next slide

- Observing
  - Pale, cyanotic, or flushed skin
  - Pedal edema
  - Sacral edema
  - Oxygen saturation, or Sp0<sub>2</sub>, reading less than 95 percent on the pulse oximeter



Signs and symptoms of breathing difficulty.

© Ray Kemp/911 Imaging

- Observing
  - Noisy breathing
    - Audible wheezing (heard without stethoscope)
    - Gurgling
    - Snoring
    - Crowing
    - Stridor
    - Coughing

continued on next slide

All Rights Reserved

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

- Auscultating
  - Lung sounds on both sides during inspiration and expiration

#### Assessment: Auscultation



6. Reassess the patient's level of distress and vital signs.

- Auscultating
  - Wheezes
    - High-pitched sounds created by air moving through narrowed air passages
  - Crackles

ALWAYS LEARNING

 Fine crackling or bubbling sound heard on inspiration and caused by fluid in alveoli or by opening of closed alveoli

continued on next slide

- Auscultating
  - Rhonchi
    - Lower-pitched sounds resembling snoring or rattling, caused by secretions in larger airways
  - Stridor

ALWAYS LEARNING

 High-pitched, upper-airway sounds indicating partial obstruction of trachea or larynx

continued on next slide

- Evaluating vital sign changes, which may include:
  - Increased or decreased pulse rate
  - Changes in breathing rate
  - Changes in breathing rhythm
  - Hypertension or hypotension

#### Patient Care

- Breathing difficulty
  - Assure adequate ventilations.
  - If breathing is inadequate, begin artificial ventilation.
  - If breathing is adequate, use a nonrebreather mask at 15 liters per minute.

#### Patient Care



4. Use settings as defined in your protocols.

#### Patient Care

- Breathing difficulty
  - Place patient in position of comfort.
  - Administer prescribed inhaler.
  - Administer continuous positive airway pressure (CPAP).

## Continuous Positive Airway Pressure (CPAP)

- Simple principles
  - Blowing oxygen or air continuously at low pressure into airway prevents alveoli from collapsing at end of exhalation.
  - Can prevent fluid shifting into alveoli from surrounding capillaries

# Continuous Positive Airway Pressure (CPAP)

- Common uses
  - Pulmonary edema
  - Drowning
  - Asthma and COPD
  - Respiratory failure in general

# Continuous Positive Airway Pressure (CPAP)

- Contraindications
  - Severely altered mental status
  - Lack of normal, spontaneous respiratory rate
  - Inability to sit up
  - Hypotension/shock

ALWAYS LEARNING

## Continuous Positive Airway Pressure (CPAP)

- Contraindications
  - Nausea and vomiting
  - Penetrating chest trauma
  - Shock

ALWAYS LEARNING

- Upper GI bleeding or recent gastric surgery
- Conditions preventing good mask seal

# Continuous Positive Airway Pressure (CPAP)

- Side effects
  - Hypotension
  - Pneumothorax
  - Increased risk of aspiration
  - Drying of corneas

All Rights Reserved

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

ALWAYS LEARNING

## Continuous Positive Airway Pressure (CPAP)

- Explain procedure to patient.
- Start with low level CPAP.

### Patient Care: Using CPAP



2. Explain the device to the patient. The mask and snug seal may initially cause the patient to feel smothered and anxious.

## Continuous Positive Airway Pressure (CPAP)

- Reassess patient's mental status, vital signs, and dyspnea level frequently.
- Raise CPAP level if no relief within a few minutes.

## Patient Care: Using CPAP




5. Reassess and monitor the patient.

## Continuous Positive Airway Pressure (CPAP)

 If patient deteriorates, remove CPAP and begin ventilating with bag mask.

## Patient Care: Using CPAP



6. Discontinue CPAP and ventilate the patient if breathing becomes inadequate.

#### **Respiratory Conditions**

# Chronic Obstructive Pulmonary Disease (COPD)

- Broad classification of chronic lung diseases
- Includes emphysema, chronic bronchitis, and black lung
- Overwhelming majority of cases are caused by cigarette smoking.


continued on next slide

ALWAYS LEARNING

# Chronic Obstructive Pulmonary Disease (COPD)

- Chronic bronchitis
  - Bronchiole lining inflamed
  - Excess mucus produced
  - Cells in bronchioles that normally clear away mucus accumulations are unable to do so

#### COPD: Chronic Bronchitis



Chronic bronchitis and emphysema are chronic obstructive pulmonary diseases.

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

All Rights Reserved

# Chronic Obstructive Pulmonary Disease (COPD)

- Emphysema
  - Alveoli walls break down.
    - Surface area for respiratory exchange is greatly reduced.
  - Lungs lose elasticity.
  - Results in air with carbon dioxide being trapped in lungs, reducing effectiveness of normal breathing

#### **Asthma**

- Chronic disease with episodic exacerbations
- During attack, small bronchioles narrow (bronchoconstriction); mucus is overproduced.
- Results in small airway passages practically closing down, severely restricting air flow

#### **Asthma**

- Airflow mainly restricted in one direction
- Inhalation
  - Expanding lungs exert outward pull, increasing diameter of airway and allowing air flow into lungs.
- Fxhalation
  - Opposite occurs and air becomes trapped in lungs.

- Abnormal accumulation of fluid in alveoli
- Patients with congestive heart failure (CHF) may experience difficulty breathing because of this.

continued on next slide

ALWAYS LEARNING

- Pressure builds up in pulmonary capillaries.
- Fluid crosses the thin barrier and accumulates in the alveoli.
- Fluid occupying lower airways makes it difficult for oxygen to reach blood.
- Patient experiences dyspnea.

- Common signs and symptoms
  - Dyspnea
  - Anxiety
  - Pale and sweaty skin
  - Tachycardia
  - Hypertension
  - Respirations are rapid and labored.
  - Low oxygen saturation

- Common signs and symptoms
  - In severe cases, crackles or sometimes wheezes may be audible.
  - Patients may cough up frothy sputum, usually white, but sometimes pinktinged.

continued on next slide

ALWAYS LEARNING

- Treatment
  - Assess for and treat inadequate breathing.
  - High-concentration oxygen
  - If possible, keep patient's legs in dependent position.
  - CPAP may be used to push fluid back out of lungs and into capillaries.

#### Think About It

- Might it be possible for a patient to have multiple respiratory disorders?
- Could a person with an underlying diagnosis of COPD also have pulmonary edema?

- Infection of one or both lungs caused by bacteria, viruses, or fungi
- Results from inhalation of certain microbes
- Microbes grow in lungs and cause inflammation.

continued on next slide

All Rights Reserved

- Signs and symptoms
  - Shortness of breath with or without exertion
  - Coughing
  - Fever and severe chills
  - Chest pain (often sharp and pleuritic)

- Signs and symptoms
  - Headache
  - Pale, sweaty skin
  - Fatigue
  - Confusion

continued on next slide

All Rights Reserved

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

- Treatment
  - Care mostly supportive
  - Assess for and treat inadequate breathing.
  - Oxygenate
  - Transport

### Spontaneous Pneumothorax

- Lung collapses without injury or other obvious cause.
- Tall, thin people, and smokers are at higher risk for this condition.

continued on next slide

ALWAYS LEARNING

#### Spontaneous Pneumothorax

- Signs and symptoms
  - Sharp, pleuritic chest pain
  - Decreased or absent lung sounds on side with injured lung
  - Shortness of breath/dyspnea on exertion
  - Low oxygen saturation, cyanosis
  - Tachycardia

ALWAYS LEARNING

#### Spontaneous Pneumothorax

- Treatment
  - Transport for definitive care, as patients frequently require chest tube.
  - Administer oxygen.
  - CPAP contraindicated

## Pulmonary Embolism

- Blockage in blood supply to lungs
- Commonly caused by deep vein thrombosis (DVT)
- Increased risk from limb immobility, local trauma, or abnormally fast blood clotting

## Pulmonary Embolism

- Signs and symptoms
  - Sharp, pleuritic chest pain
  - Shortness of breath
  - Anxiety
  - Coughing
  - Sweaty skin that is pale or cyanotic
  - Tachycardia
  - Tachypnea
  - Wheezing

## Pulmonary Embolism

- Treatment
  - Difficult to differentiate in field
  - Transport to definitive care.
  - Oxygenate.

## **Epiglottitis**

- Infection causing swelling around and above the epiglottis.
- In severe cases, swelling can cause airway obstruction.

continued on next slide

All Rights Reserved

Copyright © 2016, 2012, 2009 by Pearson Education, Inc.

## **Epiglottitis**

- Signs and symptoms
  - Sore throat, drooling, difficult swallowing
  - Preferred upright or tripod position
  - Sick appearance
  - Muffled voice
  - Stridor

ALWAYS LEARNING

continued on next slide

## **Epiglottitis**

- Treatment
  - Keep patient calm and comfortable.
  - Do not inspect throat.
  - Administer high-concentration oxygen if possible without alarming patient.
  - Transport.

## Cystic Fibrosis

- Genetic disease typically appearing in childhood
- Causes thick, sticky mucus accumulating in the lungs and digestive system
- Mucus can cause life-threatening lung infections and serious digestion problems.

continued on next slide

All Rights Reserved

## Cystic Fibrosis

- Signs and symptoms
  - Coughing with large amounts of mucus
  - Fatigue
  - Frequent occurrences of pneumonia
  - Abdominal pain and distention
  - Coughing up blood
  - Nausea
  - Weight loss

continued on next slide

## Cystic Fibrosis

- Treatment
  - Caregiver often best resource for baseline assessment of patient.
  - Caregivers can often guide treatment.
  - Assess for, and treat, inadequate breathing.
  - Transport.

## Viral Respiratory Infections

- Infection of respiratory tract
- Usually minor but can be serious, especially in patients with underlying respiratory diseases like COPD

continued on next slide


All Rights Reserved

ALWAYS LEARNING

## Viral Respiratory Infections

- Often starts with sore or scratchy throat with sneezing, runny nose, and fatigue
- Fever and chills
- Infection can spread into lungs, causing shortness of breath.
- Cough can be persistent.
  - May produce yellow or greenish sputum


## Chronic Obstructive Pulmonary Diseases Video



Click on the screenshot to view a video on the subject of chronic obstructive pulmonary diseases.

**Back to Directory** 


## Spontaneous Pneumothorax Animation



Click on the screenshot to view an animation on the subject of spontaneous pneumothorax.


**Back to Directory** 

- Metered-dose inhaler
- Provides a metered (exactly measured) inhaled dose of medication
- Most commonly prescribed for conditions causing bronchoconstriction



Prescribed Inhaler

- Before administering inhaler
  - Right patient, right time, right medication, right dose, right route
  - Check expiration date.
  - Shake inhaler vigorously.
- Patient alert enough to use inhaler
  - Use spacer device if patient has one.



3. Ensure the five "rights": 1. Right patient; 2. Right time; 3. Right medication; 4. Right dose; 5. Right route.

## Spacer Device



A spacer between the inhaler and patient makes the timing during inhaler use less critical.

- To administer inhaler:
  - Have patient exhale deeply.
  - Have patient put lips around opening.
  - Press inhaler to activate spray as patient inhales deeply.
  - Make sure patient holds breath as long as possible so medication can be absorbed.



Have the patient seal his lips around the mouthpiece and breathe deeply. Instruct the patient to hold his breath for 2 to 3 seconds if possible. Continue until the medication is gone from the chamber.

# Using a Metered Dose Asthma Inhaler and Spacer Video



Click on the screenshot to view a video on the subject of using a metered dose inhaler.

**Back to Directory** 

#### The Small-Volume Nebulizer

#### The Small-Volume Nebulizer

- Medications used in metered-dose inhalers can also be administered by a small-volume nebulizer (SVN).
- Nebulizing
  - Running oxygen or air through liquid medication
- Patient breathes vapors created.

#### The Small-Volume Nebulizer

- Produces continuous flow of aerosolized medication that can be taken in during multiple breaths over several minutes
- Gives patient greater exposure to medication

 Respiratory emergencies are common complaints for EMTs. It is important to understand the anatomy, physiology, pathophysiology, assessment, and care for patients experiencing these emergencies.

continued on next slide

ALWAYS LEARNING

 Patients with respiratory complaints (which are closely related to cardiac complaints) may exhibit inadequate breathing. Rapid respirations indicate serious conditions including hypoxia, cardiac and respiratory problems, and shock.

continued on next slide

ALWAYS LEARNING

 Very slow and shallow respirations are often the endpoint of a serious condition and are a precursor to death.

continued on next slide

 The history usually provides significant information about the patient's condition. In addition to determining a pertinent past history and medications, determine the patient's signs and symptoms with a detailed description including OPQRST and events leading up to the episode.

continued on next slide

 Important physical examination points include checking the patient's work of breathing, inspecting accessory muscle use, gathering pulse oximetry readings, assuring adequate and equal lung sounds bilaterally, examining for excess fluid (lungs, ankles, and abdomen), and gathering vital signs.

continued on next slide

ALWAYS LEARNING

#### Remember

- Determine if the patient's breathing is adequate, inadequate, or absent.
- Choose the appropriate oxygenation or ventilation therapy.

continued on next slide

 Several medications are available that may help correct a patient's difficulty in breathing.

#### Remember

- Consider whether to assist a patient with or administer respiratory medications.
  - Do I have protocols and medications that may help this patient?
  - Does the patient have a presentation and condition that may fit these protocols?

continued on next slide

ALWAYS LEARNING

#### Remember

- Consider whether to assist a patient with or administer respiratory medications.
  - Are there any contraindications or risks to using medications in my protocols?

#### Questions to Consider

- What would you expect a patient's respiratory rate to do when the patient gets hypoxic? Why?
- What would you expect a patient's pulse rate to do when the patient gets hypoxic? Why?
- List the signs of inadequate breathing.

continued on next slide

#### **Questions to Consider**

- Would you expect to assist a patient with their prescribed inhaler when they are experiencing congestive heart failure? Why or why not?
- List some differences between adult and infant/child respiratory systems.

## Critical Thinking

 A 72-year-old female complains of severe shortness of breath. Her husband notes she is confused. You note respiratory rate of 8 breaths/minute and cyanosis. Patient has a history of COPD and CHF. Discuss the treatment steps to assist this patient.